

Smooth tail index estimation

Samuel Müller^{a*}, Kaspar Rufibach^{b†}

^aSchool of Mathematics and Statistics, University of Western Australia
Crawley, WA 6009, Australia

^bDepartment of Statistics, Stanford University
Stanford, CA 94305, United States

February 2, 2008

Abstract

Both parametric distribution functions appearing in extreme value theory - the generalized extreme value distribution and the generalized Pareto distribution - have log-concave densities if the extreme value index $\gamma \in [-1, 0]$. Replacing the order statistics in tail index estimators by their corresponding quantiles from the distribution function that is based on the estimated log-concave density $\hat{f}_n$ leads to novel smooth quantile and tail index estimators. These new estimators aim at estimating the tail index especially in small samples. Acting as a smoother of the empirical distribution function, the log-concave distribution function estimator reduces estimation variability to a much greater extent than it introduces bias. As a consequence, Monte Carlo simulations demonstrate that the smoothed version of the estimators are well superior to their non-smoothed counterparts, in terms of mean squared error.

Keywords: “extreme value” theory; log-concave density estimation; negative Hill estimator; Pickands estimator; tail index estimation; small-sample performance

2000 Mathematics Subject Classifications: Primary 62G32, 62G07; Secondary 60G70

1 Introduction

It is a well-known fact that asymptotic results are in general at best approximately valid in small-sample problems, but that in the latter situation bias is often a serious issue. For

*Corresponding author. Email: mueller@maths.uwa.edu.au (S. Müller), tel: +61 (0)8 6488 7173

†K. Rufibach (kaspar.rufibach@stanford.edu) is partially supported by Swiss National Science foundation.

example in extreme value theory the small-sample bias in the estimation of the tail index is severely. We refer to [6] for a study of a number of estimators. There are only a few more articles that focus on the small-sample performance of tail-index estimators. We are aware of [2], [22] and [38]. These consider mainly tail-index estimation for heavy-tailed distributions. In this article we investigate the small-sample behavior of a new smooth tail-index estimator for thin-tailed and distributions with finite endpoint. The main aim of this article is to introduce the smoothed estimators that exploit the log-concavity of the limiting density of the exceedances or of the largest order statistics, respectively. In Section 2 we present the connection of log-concavity and extreme-value theory. Further, we show in Section 3 that replacing the empirical distribution function by the smooth estimator $\hat{F}_n$ based on the log-concave density estimator - see (2) for a proper definition of $\hat{F}_n$ - leads to novel tail-index estimators that exhibit substantially decreased mean-squared error in small-sample situations. We illustrate this finding in a simulation study in Section 4 for two settings (1) a generalized Pareto distribution and (2) a domain of attraction scenario. The paper concludes with some brief remarks in Section 5.

2 Log-concavity in extreme-value theory

2.1 Max-domain of attraction of distributions with log-concave densities.

Let $\{X_n, n \geq 1\}$ be a sequence of independent random variables with common distribution function F , such that F belongs to the max-domain of attraction of G , denoted by $F \in \mathcal{D}(G)$, i.e. there exist constants $a_n > 0$, $b_n \in \mathbb{R}$ such that for $x \in \mathbb{R}$ and

$$\begin{aligned} G(x) &= \lim_{n \rightarrow \infty} \mathbb{P} \left(a_n^{-1} \cdot [\max(X_1, \dots, X_n) + b_n] \leq x \right) \\ &= \lim_{n \rightarrow \infty} F^n(a_n \cdot x - b_n) \end{aligned}$$

which is equivalent to

$$\sup_{x \in \mathbb{R}} |F^n(a_n \cdot x - b_n) - G(x)| \rightarrow 0, \text{ as } n \rightarrow \infty.$$

From [15] it is known that $F \in \mathcal{D}(G)$ if and only if $G \in \{G_\gamma : \gamma \in \mathbb{R}\}$, where

$$G_\gamma(x) = \exp \left(-(1 + \gamma x)^{-1/\gamma} \right), \quad 1 + \gamma x > 0,$$

where G_γ is called the extreme value distribution with tail index $\gamma \in \mathbb{R}$, shift parameter 0, and scale parameter 1. Since

$$(1 + \gamma x)^{-1/\gamma} \rightarrow \exp(-x), \quad \text{for } \gamma \rightarrow 0,$$

interpret $G_0(x)$ as $\exp(-e^{-x})$. The two most common settings in the analysis of extreme values are that we either have an observed sequence of independent and identically distributed maxima, $M_{n,1}, \dots, M_{n,k}$, or upper order statistics, $X_{(n)} \geq X_{(n-1)} \geq \dots \geq X_{(n-K_n)}$, from an independent and identically distributed sample $X_1, \dots, X_n$ with $K_n = k_n < n$ or $K_n = \#\{X_i : X_i \geq u_n\}$ a random number, where u_n is some suitably chosen high threshold and $\#A$ denotes the number of elements in set A . We will focus on the latter setting and in general we assume that K_n is an intermediate sequence that is $K_n/n \rightarrow 0$ and $K_n \rightarrow \infty$ if $n \rightarrow \infty$.

2.2 The Generalized Pareto Distribution and log-concavity.

To fix notation, define for a general distribution function F the lower endpoint $\alpha(F) := \inf\{x \in \mathbb{R} : F(x) > 0\}$ and the upper endpoint $\omega(F) := \sup\{x \in \mathbb{R} : F(x) < 1\}$. The quantile function of F for $q \in [0, 1]$ is

$$F^{-1}(q) := \inf\{x \in \mathbb{R} : F(x) \leq q\}.$$

Exceedances $X_{(n)} - u, \dots, X_{(n-k_n+1)} - u$ of a high threshold $u = u_n$ or of an intermediate order statistic $u = X_{(n-k_n)}$ are typically modelled by the generalized Pareto distribution (GPD), established by Pickands (see [29]). For $\gamma \in \mathbb{R}$ and $\sigma \in (0, \infty)$ the density of the GPD is given by

$$w_{\gamma, \sigma}(x) := \sigma^{-1}(1 + \gamma x/\sigma)^{-(1+1/\gamma)}, \quad x \in [0, -\sigma/\gamma],$$

where $w_{0, \sigma}$ is again defined via continuity: $w_{0, \sigma}(x) = \sigma^{-1} \exp(-x/\sigma)$ for $x \in [0, \infty)$.

[28] showed that both parametric distribution functions appearing in extreme value theory - the generalized extreme value distribution (GEV) and the generalized Pareto distribution - have log-concave densities if the extreme value index $\gamma \in [-1, 0]$ and that all distribution functions F with log-concave density belong to the max-domain of attraction of the GEV with $\gamma \in [-1, 0]$. For any distribution function F with corresponding real-valued log-concave density function f , this latter f can be written as

$$f(x) = \exp \varphi(x), \tag{1}$$

for a concave function $\varphi : \mathbb{R} \rightarrow [-\infty, \infty)$. We will denote the class of distribution functions F having a log-concave density on its support $S = [\alpha(F), \omega(F)]$ by $\mathcal{F}_{\cap}(S)$.

2.3 The restriction $\gamma \in [-1, 0]$ and connection to bump hunting.

We are aware that the restriction of log-concave densities and of $\gamma \in [-1, 0]$, respectively, is a drawback if there is not sufficient evidence to assume that this restriction holds. On

the other hand there are often good reasons to assume that some distribution function F has all its moments finite or that its support is finite, implying $\gamma \in [-1, 0]$. Estimating the finite endpoint $\omega(F)$ of a distribution F is linked to the problem of estimating $\gamma < 0$ and its theory is well developed, see for example [18], [5], [19] and [13]. The data set of the total life span of people who died in the Netherlands, who were born between the years 1877-1881, as analyzed in [1] is a real life example that results in an estimated finite endpoint and an estimated tail index between $-1/2$ and 0. Further data sets on survival times of 208 mice exposed to radiation and on the men's 100m running times of the 1988 and 1992 Olympic Games are analyzed in [19]. By definition the distribution of the distance of two points in a closed convex set has finite support and there are many open problems regarding its limit behavior. For the current state of research we refer to [25]. A further example that naturally leads to the restriction $\gamma < 0$ is the estimation of the efficiency frontier in economics (see [14]). In practical applications, $\gamma = -1/2$ is often seen as natural lower bound, e.g. p. 62 in [24] or [13].

In this context, we would like to point out [30], who proposes a multiscale procedure to identify collections of intervals based on an i.i.d. sample where a density is either log-concave or log-convex. The chosen multiscale approach ensures that the claims (i.e. log-concavity or log-convexity) holds for all intervals in the collections simultaneously. Additionally, it is shown that the method asymptotically keeps the level. This offers a way to “pre-assess” whether γ actually is in $[-1, 0]$: For a chosen significance level $\alpha \in (0, 1)$, apply this novel bump hunting method to either the sample of exceedances or the whole sample of observations. In the first case, if the collection of intervals claiming log-convexity is empty and the collection of intervals claiming log-concavity at best contains an interval (almost) spanning the whole range of exceedances, then we could claim with asymptotic probability $1 - \alpha$ that indeed the observations we are looking at stem from a distribution $F \in \mathcal{F}_{\bar{\cap}} \subset \mathcal{D}(G_{\gamma})$, $\gamma \in [-1, 0]$.

In the second case, we get an upper bound $m \geq k$ for the number of upper order statistics we should take into account. Just define m such that $X_{(n-m+1)}$ is the left-most endpoint of all intervals whereon the bump-hunting method claims log-concavity of the underlying density.

In both cases, the new tail index estimation procedures as presented in Section 3 is adequate.

3 Tail index estimation

The estimation of γ is besides the related high quantile estimation the most important problem in univariate extreme value theory and there exist a vast number of different

approaches. For example the Hill estimator ([20]), the maximum likelihood estimator ([3], [18], [35], [36], [37]), the moment estimator ([7]), the (iterated) negative Hill estimator also known as Falk's estimator ([11], [12], [27]), the (generalized) Pickands estimator ([29], [8], [34]), estimators based on near extremes ([26]), the weighted least squares estimator ([23]), probability weighted moments ([21]), and many more. All these estimators are based on an intermediate sequence of upper order statistics and it is well known (see e.g. [17]) that a major drawback of such estimators is their discrete character. Using kernel-type estimators is one possibility to overcome this deficiency. We refer to [4] for the smoothed Hill estimator in case $\gamma > 0$ and to [17] for general $\gamma \in \mathbb{R}$. Our alternative is to take advantage of the distribution function $\widehat{F}_n$ based on the log-concave density estimator $\widehat{f}_n$, which is possible if the true γ is in the restricted interval $[-1, 0]$.

3.1 Motivation of new estimators.

For an i.i.d. sample $X_1, \dots, X_n$ where X_i has a log-concave density function as introduced in (1), let $\mathbb{F}_n$ be the empirical distribution function and

$$\widehat{F}_n(x) := \int_{-\infty}^x \widehat{f}_n(t) dt \quad (2)$$

be the smoothed distribution function based on the log-concave density estimator $\widehat{f}_n = \exp \widehat{\varphi}_n$. A proper definition and properties of $\widehat{f}_n$ are given in [30] and [10]. We only mention one special feature: the estimator $\exp \widehat{\varphi}_n$ of the log-density is a piecewise linear function with knots only at some of the observations points $X_1, \dots, X_n$ and $\widehat{\varphi}_n = -\infty$ on $\mathbb{R} \setminus [X_1, X_n]$. How to actually compute $\widehat{f}_n$ is detailed in [31] and [9].

In [30] the following theorem is proven.

Theorem 3.1. *Let $X_1, \dots, X_n$ be an i.i.d. sample stemming from a distribution with log-concave density f such that $f = \exp \varphi$ is Hölder-continuous for an exponent $\beta \in (1, 2]$ and constant $L > 0$. Furthermore, $\varphi'(x) - \varphi'(y) \geq C(y - x)$ for $C > 0$ and $A \leq x < y \leq B$. Then, as $n \rightarrow \infty$,*

$$\max_{t \in T_n} |\mathbb{F}_n - \widehat{F}_n|(t) = o_p(n^{-1/2}),$$

where $T_n \rightarrow T := [A, B]$. Furthermore, $\widehat{F}_n(X_{(1)}) = 0$ and $\widehat{F}_n(X_{(n)}) = 1$.

This theorem implies that $\widehat{F}_n$ is essentially equivalent to $\mathbb{F}_n$, but as the integral of a piecewise exponential function very smooth. These properties turn out to be highly convenient in the estimation of the extreme value tail index γ . The smoothness of $\widehat{F}_n$ reduces the variance not only considerably in the estimation of γ but even for the estimation of quantiles of the generalized Pareto distribution, as is shown in Section 4.

Many well-known tail index estimators are based on a selection of log-spacings of the sample, see also [34]. The key idea is now simply to replace the order statistics (or quantiles of the empirical distribution function) $X_{(i)} = \mathbb{F}_n^{-1}(i/n)$, $i = 1, \dots, n$ in these log-spacings by quantiles received via $\hat{F}_n$. This yields modified versions of the uniformly minimum variance unbiased estimator from Falk ([11], [12]) for the case of a known endpoint, the negative Hill estimator as defined in [12] and Pickands' estimator ([29]) for the case of a unknown endpoint. We will denote these new estimators as “smoothed estimators”. We choose the first two estimators because of their outstanding performance for $\gamma < 0$ and $\gamma < -1/2$, respectively. On the other hand it is well known that Pickands' estimator is not efficient and in addition it is able to estimate any $\gamma \in \mathbb{R}$, thus the comparison of the original to the smoothed version is not entirely fair. However, Pickands' estimator serves as the building block for much more efficient generalized Pickands' estimators that are more general linear combinations of log-spacings of order statistics ([34]).

3.2 Global and tail behavior.

Extreme value theory is, as the name suggests, tail focused. Hence, the behavior of the conditional distribution $F_{X|X>u}$, where $u \rightarrow \omega(F)$, dominates the limit results. On the other hand, the log-concavity of $f = F'$ is a strong assumption on the entire shape of the distribution function F . If this strong assumption holds then the smoothing of the tail index estimators should be based on $\hat{F}_n$. Since tail index estimators use information of the upper tail of $\mathbb{F}_n$ it was sufficient that the upper tail of F had log-concave density. Therefore, we investigate two settings in our simulation study in Section 4. (1) $X_1, \dots, X_n$ iid $\mathcal{L}(X) \in \mathcal{F}_{\bar{\cap}}(S)$. Here the data is sharpened by $\hat{F}_n$. (2) $X_1, \dots, X_n$ iid $\mathcal{L}(X) \in \mathcal{D}(G_\gamma; \gamma \in [-1, 0])$. In this more general situation F is not necessarily in $\mathcal{F}_{\bar{\cap}}(S)$. Before the smooth tail index estimators are computed, the range of log-concavity of f has to be determined; either using bump-hunting or by imposing some assumptions on F . Here the data is sharpened by $\hat{F}_m$, where the smoothed distribution function is based on the $m > K_n$ largest order statistics and m is such that $F_{X|X>X_{n-m}} \in \mathcal{F}_{\bar{\cap}}(S \cap [X_{n-m}, \infty))$.

3.3 Smooth tail index estimators.

First, let us pin down some notation. Suppose we are given a sample $X_1, \dots, X_n$ from a GPD wherefrom we know that $\gamma \in [-1, 0]$ and with empirical distribution function $\mathbb{F}_n$. Denote the order statistics by $X_{(1)}, \dots, X_{(n)}$. For such a fixed sample, define for $k = 4, \dots, n$ and $H \in \{\mathbb{F}_n, \hat{F}_n\}$:

$$\hat{\gamma}_{\text{Pick}}^k(H) = \frac{1}{\log 2} \log \left(\frac{H^{-1}((n - r_k(H) + 1)/n) - H^{-1}((n - 2r_k(H) + 1)/n)}{H^{-1}((n - 2r_k(H) + 1)/n) - H^{-1}((n - 4r_k(H) + 1)/n)} \right),$$

where

$$r_k(H) = \begin{cases} \lfloor k/4 \rfloor & \text{if } H = \mathbb{F}_n, \\ k/4 & \text{if } H = \hat{F}_n, \end{cases}$$

with $\lfloor m \rfloor := \max\{n \in \mathbb{N}_0 : n \leq m\}$. This construction not only exploits the superiority of the quantile estimates based on the smooth function $\hat{F}_n$, but also avoids “rounding bias”. Using the inverse of a continuous distribution function, quantiles do not coincide for four consecutive k ’s (order statistics), as it is the case for Pickands’ original estimate.

To generalize the estimators in [11] and [12], no discrimination regarding continuity of H is necessary. For $H \in \{\mathbb{F}_n, \hat{F}_n\}$ let

$$\begin{aligned} \hat{\gamma}_{\text{Falk}}^k(H) &= \frac{1}{k-1} \sum_{j=2}^k \log\left(\frac{X_{(n)} - H^{-1}((n-j+1)/n)}{X_{(n)} - H^{-1}((n-k)/n)}\right), \quad k = 3, \dots, n-1 \\ \hat{\gamma}_{\text{MVUE}}^k(H) &= \frac{1}{k} \sum_{j=1}^k \log\left(\frac{\omega(F) - H^{-1}((n-j+1)/n)}{\omega(F) - H^{-1}((n-k)/n)}\right), \quad k = 2, \dots, n-1 \end{aligned}$$

where F is the true distribution function of the X_i ’s. Note that $\hat{\gamma}_{\text{MVUE}}^k(H)$ is only consistent if $\gamma \in [-1, 0)$. The chosen terminology reminds of the fact that when choosing $H = \mathbb{F}_n$, the above estimators boil down to Pickands’, Falk’s, and Falk’s MVUE estimator as discussed at the beginning of this section.

The new, smooth tail index estimators are now simply $\hat{\gamma}_{\text{Pick}}^k(\hat{F}_n)$, $\hat{\gamma}_{\text{Falk}}^k(\hat{F}_n)$, and $\hat{\gamma}_{\text{MVUE}}^k(\hat{F}_n)$. Figure 1 displays Hill plots for two GPD pseudo-random samples, i.e. plots of the estimators versus the number of order statistics k , for the smoothed and unsmoothed versions for $n = 64$ and $\gamma \in \{-0.1, -0.75\}$.

The smoothed estimators behave much more stable as a function of k and it is especially noteworthy that for the two generated data sets all three smoothed estimators $\hat{\gamma}_{\text{Pick}}^k(\hat{F}_n)$, $\hat{\gamma}_{\text{Falk}}^k(\hat{F}_n)$, $\hat{\gamma}_{\text{MVUE}}^k(\hat{F}_n) \in [-1, 0]$ for every k . By construction, $\hat{\gamma}_{\text{Pick}} : \mathbb{R}^n \rightarrow \mathbb{R}$ and $\hat{\gamma}_{\text{Falk}}, \hat{\gamma}_{\text{MVUE}} : \mathbb{R}^n \rightarrow [-\infty, 0)$ which means that non permissible estimates outside the interval $[-1, 0]$ potentially occur. However, due to consistency of $\mathbb{F}_n$ and $\hat{F}_n$, this is asymptotically negligible. If in practice $\hat{\gamma} \notin [-1, 0]$ then, a truncation to its closest boundary value is recommendable if it is known that $\gamma \in [-1, 0]$.

It is beyond the scope and not the primary goal of this article to discuss the asymptotic behavior of the smoothed tail index estimators.




Figure 1: Hill plots for $n = 64$ and $\gamma = -0.1$ (plots in the upper row) and $\gamma = -0.75$ (plots in the lower row), smoothed (—) and original (---) versions.

3.4 Further shape constraints.

Straightforward computation yields the following lemma.

Lemma 3.1. *The density $w_{\gamma,\sigma}$ has the following qualitative properties which do not depend on the value of the scale parameter σ :*

<i>property</i>	<i>parameter range</i>
<i>convex non-decreasing</i>	$\gamma \leq -1$
<i>concave non-increasing</i>	$\gamma \in [-1, -1/2]$
<i>convex non-increasing</i>	$\gamma \geq -1/2$
<i>log-concave</i>	$\gamma \in [-1, 0]$
<i>log-convex</i>	$\gamma \in (-\infty, -1] \cup [0, \infty)$

Table 1: Form of $w_{\gamma,\sigma}$ as determined by the tail index γ .

Since e.g. a density estimator $\tilde{f}_n$ in the class of convex decreasing densities is available, see [16], the latter lemma raises the possibility to define smooth estimators of the tail index for other ranges of γ . However, especially in the latter case, slight modifications may be necessary, since $\tilde{F}_n(X_{(n)}) \neq 1$. Furthermore, maximum likelihood estimators under other constraints may not be as smooth as $\hat{F}_n$, since e.g. the estimator of a convex decreasing density is piecewise linear, whereas for log-concave densities this form appears for the estimator of the log-density.

3.5 Computational details.

These new smoothed estimators are made available in the R-package `smoohtail` (see [33]). This latter package depends on the package `logcondens` ([32]), which offers two algorithms for the (weighted) estimation of an arbitrary log-concave density from an i.i.d. sample of observations. Both these packages are available from CRAN.

4 Simulations

4.1 Estimation of quantiles.

The computation of the non-smoothed estimators $\hat{\gamma}_{\text{Falk}}^k(\mathbb{F}_n)$, $\hat{\gamma}_{\text{Pick}}^k(\mathbb{F}_n)$ and $\hat{\gamma}_{\text{MVUE}}^k(\mathbb{F}_n)$ heavily relies on the order statistics $X_{(i)}, i = 1, \dots, n$. But, these simply estimate the quantiles $W_{\gamma, \sigma}^{-1}(i/n)$ of the distribution whereof we want to estimate the tail index. Therefore, the accuracy of the tail index estimators is closely connected to the ability of estimating these quantiles $W_{\gamma, \sigma}^{-1}(i/n)$. To illustrate the superiority of log-concave quantile estimation over simply taking order statistics, we calculated the relative efficiency of these two estimators. Since results were similar over an extended range of σ 's, we concentrate on the case $\sigma = 1$.

To fix notation, define the log-concave estimate $\hat{X}_{(i)} = \hat{F}_n^{-1}(i/n)$ of an order statistic $X_{(i)}$, for $i = 1, \dots, n$. Let $q_{(i)}$ denote either $\hat{X}_{(i)}$ or $X_{(i)}$, then its estimated variance and bias with respect to the i/n -quantile of a GPD $W_{\gamma, 1}$ for a fixed $\gamma \in [-1, 0]$ given simulated $q_{(i), j}$ that are based on M generated samples $X_{(1), j}, \dots, X_{(n), j}, j = 1, \dots, M$ of size n drawn from $W_{\gamma, 1}$ is defined as follows:

$$\begin{aligned} \widehat{\text{Var}}(q_{(i)}, M) &:= (M-1)^{-1} \sum_{j=1}^M \left(q_{(i), j} - (1/M) \sum_{j=1}^M q_{(i), j} \right)^2 \\ \widehat{\text{Bias}}(q_{(i)}, \gamma, M) &:= M^{-1} \left(\sum_{j=1}^M q_{(i), j} \right) - W_{\gamma, 1}^{-1}(i/n). \end{aligned}$$

The relative efficiency $\rho_{\gamma, n, M}(k)$ of log-concave quantile estimation to quantile estimation

based on order statistics is then

$$\rho_{\gamma,n,M}(k) = \frac{[\widehat{\text{Bias}}(\widehat{X}_{(k)}, \gamma, M)]^2 + \widehat{\text{Var}}(\widehat{X}_{(k)}, M)}{[\widehat{\text{Bias}}(X_{(k)}, \gamma, M)]^2 + \widehat{\text{Var}}(X_{(k)}, M)}.$$

Figure 2 details $\rho_{\gamma,32,1000}(k)$ for $\gamma \in \{-1, -0.75, -0.5, -0.25, 0\}$ as a function of k . Relative efficiencies smaller than 1 are in favor of the log-concave quantile estimation and indicate its superiority. The use of the log-concave density estimator for the estimation of quantiles substantially reduces the variance of the estimation, due to its smoothing property detailed in Theorem 3.1. This transfers to a reduced MSE, uniformly in γ and k , as is detailed in Figure 2.




Figure 2: Relative MSE in quantile estimation for $n = 32$.

4.2 Smoothed versus unsmoothed tail index estimators.

To assess the effect of smoothing the tail index estimators, we perform a simulation study for two settings. In Setting 1, we draw $M = 1000$ samples of size $n_1 = 64$ from a GPD with $\sigma = 1$ and the extreme value tail index $\gamma \in \{-1, -0.75, -0.5, -1/3, -0.25, -0.1\}$. For every k , the log-concave density is estimated based on the full sample $X_{(1)}, \dots, X_{(n)}$. Setting 2 consists of $M = 1000$ samples from a $\beta(\theta_1, \theta_2)$ -distribution having density

$$f_{\theta_1, \theta_2}(x) = \frac{\Gamma(\theta_1 + \theta_2)}{\Gamma(\theta_1)\Gamma(\theta_2)} x^{\theta_1-1} (1-x)^{\theta_2-1}, \quad x \in [0, 1], \quad \theta_1, \theta_2 > 0. \quad (3)$$

The upper tail of the β -distribution is dominated by θ_2 , since for $x \uparrow 1$ we have that $f_{\theta_1, \theta_2}(x) = c(\theta_1, \theta_2) \cdot w_{-\theta_2^{-1}, \theta_2^{-1}}(x)[1 + o(1)]$. Thus, $\beta(\theta_1, \theta_2) \in \mathcal{D}(G_{-1/\theta_2})$. We fix $\theta_1 = 1/2$ and thus only the upper tail of $f_{0.5, \theta_2}$ is log-concave. For the simulations, we choose $\theta_2 = -\gamma^{-1} \in \{1, 4/3, 2, 3, 4, 10\}$ and $n_2 = 128$ which equals deliberately $2n_1$ in order to underline the difference between the two settings. Here the log-concave density estimator is based on the largest $n_2/2 = 64$ order statistics. In both settings we present results for a single sample size only because the results for $n_1, n_2/2 \in \{32, 128, 256, 512\}$ are very similar.

Setting 1 represents the “ideal” of observing pure “peak over threshold” data

$$X_1, \dots, X_n \text{ iid } \mathcal{L}(X) = \text{GPD} \in \mathcal{F}_{\cap}(S).$$

Alternatively, Setting 2 stands for the more general situation

$$X_1, \dots, X_n \text{ iid } \mathcal{L}(X) \in \mathcal{D}(G_\gamma; \gamma \in [-1, 0]).$$

In the latter case, the well known problem of the tradeoff between bias and variance dominates the optimal choice of k whereas in Setting 1, k can be chosen as large as possible since we are considering the “perfect” model.

4.3 Simulation results.

We compute relative efficiencies for the estimation of the tail index as for quantile estimation. For Setting 1, results are displayed in Figure 3. If one knows that $\gamma \in [-1, 0]$, then clearly using the smoothed estimator is most worthwhile for Pickands’ estimator. However, also the other two estimators are substantially improved, with highest gain in terms of MSE for small k ’s.

Figure 4 sheds light on the bias-variance trade-off in Setting 2 for $\theta_2 = 3$; results for other choices of θ_2 were absolutely similar and therefore omitted. The variance in estimation of the tail index is dramatically reduced for the smoothed Pickands’ estimate. The plot of the bias against k confirms that all estimators are biased as expected, since the data is generated by (3) and not by the GPD with $\gamma \in [-1, 0]$. Especially for Pickands’ estimator, the bias is a much smoother function of k for the smoothed estimators than this is the case for the original estimator. Figure 5 shows the computed relative efficiencies for the estimation of the tail index for Setting 2. The results for $\hat{\gamma}_{\text{Pick}}$ and $\hat{\gamma}_{\text{Falk}}$ are similar to those in Setting 1, yielding the most substantial improvement for Pickands’ estimator. On the other hand the efficiency of the smoothed and original $\hat{\gamma}_{\text{MVUE}}$ is almost 1, independent of k and the choice of θ_2 .




Figure 3: Relative MSE for tail index estimation for peaks over threshold data, $n = 64$.

5 Conclusions

In this article we showed that for the class of distributions $F \in \mathcal{D}(G_\gamma)$ having log-concave densities or log-concave conditional densities given that the observations exceed some threshold u , respectively, the smoothing of the empirical distribution function by the corresponding log-concave density estimator leads to improved quantile estimation uniformly in γ and the quantile to be estimated and to more efficient tail index estimators if it is known that $\gamma \in [-1, 0]$. The reduction of the mean squared error of the smoothed tail index estimator is not surprisingly most substantial for Pickands' estimator, given the known poor efficiency of the latter. The estimator $\hat{F}_n$ smoothes the empirical distribution function based on the global property of the log-concavity of the density of the underlying distribution function F . We showed that if such a global property is present then it is of great value for the estimation of a tail property such as the extreme value index γ . Of course the price to be paid is assuming that γ is restricted to the narrow interval $[-1, 0]$ and it makes only sense to use the presented smoothed estimators if there is sufficient reason for that assumption. However, the novel bump hunting method in [30] is a possible tool for testing if a log-concavity assumption holds which automatically causes the desired restriction on γ .




Figure 4: Domain of attraction scenario: $n = 128$, $\theta_1 = 1/2$, $\theta_2 = 3$, smoothed (—) and original (---) versions.

We want to stress the fact that the Hill plots in Section 3 and the results in our simulation study are not based on an optimal choice of the setting parameters and are reproducible for a wide range of parameter values for the $\beta(\theta_1, \theta_2)$ -distribution unless the upper tail of the density is log-concave. Our simulation study can be reproduced using the R-package `smoothtail` (see [33]).

5.1 Further research.

The results in this article raise many new open questions and problems. We only mention two. At first it is challenging to prove asymptotic normality for the smoothed estimators. Then, by fitting the generalized Pareto density to $\hat{f}_n$ or $\log w_{\gamma, \sigma}$ to $\hat{\varphi}_n$, respectively, opens the door to construct novel estimators for the tail index that are worthwhile to be investigated in the future.

6 Acknowledgments.

This research was initiated while both authors were working at the Institute of Mathematical Statistics and Actuarial Science at the University of Bern. K. Rufibach was partially supported by Swiss National Science foundation.




Figure 5: Relative MSE in domain of attraction scenario, $n = 128$. θ_2 equals 10, 4 and 3 in the top row and 2, $4/3$ and 1 in the bottom row.

References

- [1] Aarssen, K., de Haan, L., 1994. On the maximal life span of humans. *Mathematical Population Studies*, 4(4) 259-281.
- [2] Brazauskas, V., Serfling, R., 2001. Small sample performance of robust estimators of tail parameters for Pareto and exponential models. *J. Statist. Comput. Simulation*, 70(1) 1-19.
- [3] Coles, S.G., Dixon, M.J., 1999. Likelihood-based inference for extreme value models. *Extremes*, 2(1) 5-23.
- [4] Csörgő, S., Deheuvels, P., Mason, D., 1985. Kernel estimates of the tail index of a distribution. *Ann. Statist.*, 13(3) 1050-1077.

- [5] Csörgő, S., Mason, D.M., 1989. Simple estimators of the endpoint of a distribution. In J. Hüsler & R.-D. Reiss (Eds.), *Extreme Value Theory, Lecture Notes in Statistics*, New York. 132-147.
- [6] Dacorogna, M., Müller, U., Pictet, O., de Vries, C., The distribution of extremal foreign exchange rate returns in extremely large data sets. Discussion Paper 95-70. (Tinbergen Institute, Netherlands, 1995).
- [7] Dekkers, A.L.M., de Haan, L., Einmahl, J.H.J., 1989. A moment estimator for the index of an extreme-value distribution. *Ann. Statist.*, 17(4) 1833-1855.
- [8] Drees, H., 1995. Refined Pickands estimator of the extreme value index. *Ann. Statist.*, 23(6) 2059-2080.
- [9] Dümbgen L., Hüsler A., Rufibach K., Active Set and EM Algorithms for Log-Concave Densities Based on Complete and Censored Data. Preprint. (Dept. of Mathematical Statistics, University of Bern, Switzerland, 2006).
- [10] Dümbgen L., Rufibach K., Maximum likelihood estimation of a log-concave density and its distribution function: basic properties and uniform consistency. Preprint. (Dept. of Mathematical Statistics, University of Bern, Switzerland, 2006).
- [11] Falk, M., 1994. Extreme quantile estimation in δ -neighborhoods of generalized Pareto distributions. *Statist. Probab. Lett.*, 20(1) 9-21.
- [12] Falk, M., 1995. Some best parameter estimates for distributions with finite endpoint. *Statistics*, 27(1-2) 115-125.
- [13] Ferreira, A., de Haan, L., Peng, L., 2003. On optimising the estimation of high quantiles of a probability distribution. *Statistics*, 37(5) 401-434.
- [14] Farrell, M.J., 1957. The measurement of productive efficiency. *J. Roy. Statist. Soc. Ser. A*, 120(3) 253-281.
- [15] Gnedenko, B.V., 1943. Sur la distribution limite du terme maximum d'une série aléatoire. *Ann. of Math. (2)*, 44(3) 423-453.
- [16] Groeneboom, P., Jongbloed, G., Wellner, J.A., 2001. Estimation of a convex function: characterization and asymptotic theory. *Ann. Statist.*, 29(6) 1653-1698.
- [17] Groeneboom, P., Lopuhaä, H.P., de Wolf, P.P., 2003. Kernel-type estimators for the extreme value index. *Ann. Statist.*, 31(6) 1956-1995.
- [18] Hall, P., 1982. On estimating the endpoint of a distribution. *Ann. Statist.*, 10(2) 556-568.
- [19] Hall, P., Wang, J.Z., 1999. Estimating the end-point of a probability distribution using minimum-distance methods. *Bernoulli*, 5(1) 177-189.

- [20] Hill, B.M., 1975. A simple general approach to inference about the tail of a distribution. *Ann. Statist.*, 3(5) 1163-1174.
- [21] Hosking, J.R.M., Wallis, J.R., Wood, E.F., 1985. Estimation of the generalized extreme-value distribution by the method of probability-weighted moments. *Technometrics*, 27(3) 251-261.
- [22] Huisman, R., Koedijk, K.G., Kool, C.J.M., Palm, F., 2001. Tail-index estimates in small samples. *J. Bus. Econom. Statist.*, 19(2) 208-216.
- [23] Hüsler, J., Li, D., Müller, S., 2006. Weighted least squares estimation of the extreme value index. *Statist. Probab. Lett.*, 76(9) 920-930.
- [24] Kotz, S., Nadarajah, S., 2000. *Extreme value distributions. Theory and applications.* Imperial College Press, London.
- [25] Mayer, M., Molchanov, I., Limit theorems for the diameter of a random sample in the unit ball. Technical Report Nr 59. (Dept. of Mathematical Statistics, University of Bern, Switzerland, 2006).
- [26] Müller, S., 2003. Tail estimation based on numbers of near m -extremes. *Methodol. Comput. Appl. Probab.*, 5(2) 197-210.
- [27] Müller, S., Hüsler, J., 2005. Iterative estimation of the extreme value index. *Methodol. Comput. Appl. Probab.*, 7(2) 139-148.
- [28] Müller, S., Rüfibaich, K., On the max-domain of attraction of distributions with log-concave densities. Submitted.
- [29] Pickands, J., 1975. Statistical inference using extreme order statistics. *Ann. Statist.*, 3(3) 119-131.
- [30] Rüfibaich, K., Log-Concave Density Estimation and Bump Hunting for i.i.d. Observations. Dissertation. (Universities of Bern and Göttingen, 2006). Available on http://www.stub.unibe.ch/download/eldiss/06rufibach_k.pdf.
- [31] Rüfibaich, K., 2007. Computing maximum likelihood estimators of a log-concave density function. *J. Statist. Comput. Simul.*, to appear.
- [32] Rüfibaich, K., Dümbgen, L., 2006. logcondens: Estimate a log-concave probability density from iid observations. R package version 1.2.
- [33] Rüfibaich, K., Müller, S., 2006. smoothtail: Smooth estimation of GPD shape parameter. R package version 1.1.
- [34] Segers, J., 2005. Generalized Pickands estimators for the extreme value index. *J. Statist. Plann. Inference*, 128(2) 381-396.

- [35] Smith, R.L., 1985. Maximum likelihood estimation in a class of nonregular cases. *Biometrika*, 72(1) 67-90.
- [36] Smith, R.L., 1987. Estimating tails of probability distributions. *Ann. Statist.*, 15(3) 1174-1207.
- [37] Smith, R.L, Weissman, I., 1985. Maximum likelihood estimation of the lower tail of a probability distribution. *J. Roy. Statist. Soc. Ser. B*, 47(2) 285-298.
- [38] Wagner, N., Marsh, T.A., 2004. Tail index estimation in small samples. Simulation results for independent and ARCH-type financial return models. *Statist. Papers*, 45(4) 545-561.